



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 302 389 B1**

12

EUROPEAN PATENT SPECIFICATION

- (45) Date of publication of patent specification: 22.12.93 (51) Int. Cl.⁵: **C07D 213/36, C07D 213/74, C07D 215/12, C07D 277/38, C07D 213/71, C07D 277/28, C07D 277/32, C07D 213/61, A01N 43/40, A01N 43/42, A01N 43/78**
- (21) Application number: 88112210.5
- (22) Date of filing: 28.07.88
- Divisional application 92111470.8 filed on 28/07/88.

(54) Alpha-unsaturated amines, their production and use.

- (30) Priority: 01.08.87 JP 192793/87
13.10.87 JP 258856/87
26.01.88 JP 16259/88
17.03.88 JP 64885/88
- (43) Date of publication of application:
08.02.89 Bulletin 89/06
- (45) Publication of the grant of the patent:
22.12.93 Bulletin 93/51
- (54) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE
- (56) References cited:
EP-A- 0 002 930 EP-A- 0 092 647
EP-A- 0 164 040 EP-A- 0 254 859
DE-A- 3 232 462 DE-A- 3 343 884

- (73) Proprietor: Takeda Chemical Industries, Ltd.
1-1, Doshomachi 4-chome
Chuo-ku, OSAKA(JP)
- (72) Inventor: Minamida, Isao
5-91, Fushimidai 2-chome
Inagawa-cho
Kawabe-gun Hyogo 666-02(JP)
Inventor: Iwanaga, Koichi
8-7, Midorigaoka 2-chome
Ikeda Osaka 563(JP)
Inventor: Okauchi, Tetsuo
10-11, Tsutsumi-cho
Hirakata Osaka 573(JP)
- (74) Representative: von Kreisler, Alek,
Dipl.-Chem. et al
Patentanwälte
von Kreisler-Selting-Werner
Postfach 10 22 41
D-50462 Köln (DE)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

CHEMICAL ABSTRACTS, vol. 94, no. 9, 2nd March 1981, page 637, column 1, abstract-no. 64642], Columbus, Ohio, US; V.J. RAM: "Organosulfur compounds as potential pesticides"

JOURNAL OF MEDICINAL CHEMISTRY, vol. 27, 1984, pages 849-857; J. YANAGISAWA et al.: "Histamine H2 receptor antagonists. 1. Synthesis of N-cyano and N-carbamoyl amidine derivatives and their biological activities"

PATENT ABSTRACTS OF JAPAN, vol. 11, no. 369 (C-461)(2816), 2nd December 1987


Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. An α -unsaturated amine of the formula:

5

10




15

wherein:

20

one of X^1 and X^2 is an electron-attracting group and the other is a hydrogen atom or an electron-attracting group, wherein the said electron-attracting group is cyano, nitro, C_1-4 alkoxy carbonyl, carboxyl, C_6-10 aryloxy-carbonyl, heterocycleoxycarbonyl, C_1-4 alkylsulfonyl which may be substituted with halogen, aminosulfonyl, di- C_1-4 alkoxyphosphoryl, C_1-4 alkanoyl which may be substituted with halogen, C_1-4 alkylsulfonylthiocarbamoyl, carbamoyl or halogen, or X^1 and X^2 together with the carbon atom to which they are attached form a ring of the formula:


25



30

R^1 is a group of the formula:

35



in which:

40

R^3 is hydrogen, C_1-20 alkyl, C_6-10 aryl, C_7-9 aralkyl, heterocycle, C_1-4 alkanoyl, C_6-10 aryl-carbonyl, C_1-4 alkoxy-carbonyl, C_6-10 aryloxy-carbonyl, heterocycleoxycarbonyl, C_6-10 arylsulfonyl, C_1-4 alkylsulfonyl, di- C_1-4 alkoxyphosphoryl, C_1-4 alkoxy, hydroxy, amino, di- C_1-4 alkylamino, C_1-4 alkanoylamino, C_1-4 alkoxy-carbonylamino, C_1-4 alkylsulfonylamino, di- C_1-4 alkoxyphosphorylamino, C_7-9 aralkyloxy or C_1-4 alkoxy-carbonyl- C_1-4 alkyl; and


45

R^4 is hydrogen, C_1-20 alkyl, C_3-6 cycloalkyl, C_2-6 -alkenyl, C_3-6 cycloalkenyl or C_2-6 alkynyl, wherein each of the radicals defined for R^4 except for hydrogen may optionally be substituted by 1 to 3 substituents selected from the group consisting of hydroxy, C_1-4 alkoxy, halogen, di- C_1-4 alkylamino, C_1-4 alkylthio, C_1-3 alkanoylamino, C_1-4 alkylsulfonylamino, tri- C_1-4 alkylsilyl, pyridyl and thiazolyl, and each of the pyridyl and thiazolyl may further be substituted by halogen, or

50


R^3 and R^4 together with the adjacent nitrogen atom constitute a cyclic amino group of the formula:

55



R^2 is (1) hydrogen, (2) a group attached through a carbon atom selected from the class consisting of C_1-4 alkanoyl, C_1-20 alkyl, C_2-6 alkenyl, C_3-6 cycloalkyl, C_6-10 aryl, C_7-9 aralkyl and 3- or 4-

pyridyl, the said group attached through a carbon atom being optionally substituted by 1 to 3 substituents selected from the class consisting of C₁₋₄ alkylthio, C₁₋₄ alkoxy, mono- or di-C₁₋₄ alkylamino, C₁₋₄ alkoxy-carbonyl, C₁₋₄ alkylsulfonyl, halogen and C₁₋₄ alkanoyl, (3) a group attached through an oxygen atom selected from the class consisting of C₁₋₄ alkoxy, C₃₋₆ cycloalkoxy, C₂₋₄ alkenyloxy, C₃₋₆ cycloalkenyloxy, ethynyloxy, C₆₋₁₀ aryloxy, thienyloxy and hydroxy, the said group attached through an oxygen atom being optionally substituted by 1 to 3 substituents selected from the class consisting of halogen and phenyl, or (4) a group attached through a nitrogen atom of the formula:



wherein

R³ and R⁴ have the meanings given above;

n is an integer of 0, 1 or 2;

A^o is heterocycle;

wherein the heterocycle in the said heterocycle carbonyl for X¹ and X², the said heterocycle for R³, the heterocycle in the said heterocycleoxycarbonyl for R³, and the said heterocycle for A^o are a member selected from the class consisting of thienyl, furyl, pyrrolyl, pyridyl, oxazolyl, thiazolyl, pyrazolyl, imidazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, N-oxidopyridyl, pyrimidinyl, N-oxidopyrimidinyl, pyridazinyl, pyrazinyl, N-oxidopyridazinyl, benzofuryl, benzothiazolyl, benzoxazolyl, triazinyl, oxotriazinyl, tetrazo[1,5-b]-pyridazinyl, triazolo[4,5-b]pyridazinyl, oxoimidazinyl, dioxotriazinyl, pyrrolidinyl, piperidinyl, pyranyl, thiopyranyl, 1,4-oxazinyl, morpholinyl, 1,4-thiazinyl, 1,3-thiazinyl, piperazinyl, benzimidazolyl, quinolyl, isoquinolyl, indolizinyll, quinolizinyll, 1,8-naphthyridinyl, purinyl, pteridinyl, dibenzofuranyl, carbazolyl, acridinyl, phenanthridinyl, phenazinyl, phenothiazinyl and phenoxazinyl, the said heterocycle being optionally substituted by 1 to 5 substituents selected from the group consisting of:

- (i) C₁₋₄ alkyl,
- (ii) C₃₋₆ cycloalkyl,
- (iii) C₆₋₁₀ aryl,
- (iv) C₁₋₄ alkoxy,
- (v) C₃₋₆ cycloalkyloxy,
- (vi) C₆₋₁₀ aryloxy,
- (vii) C₇₋₁₂ aralkyloxy
- (viii) C₁₋₄ alkylthio,
- (ix) C₃₋₆ cycloalkylthio,
- (x) C₆₋₁₀ arylthio,
- (xi) C₇₋₁₂ aralkylthio,
- (xii) mono-C₁₋₄ alkylamino,
- (xiii) di-C₁₋₄ alkylamino,
- (xiv) C₃₋₆ cycloalkylamino,
- (xv) C₆₋₁₀ arylamino,
- (xvi) C₇₋₁₂ aralkylamino,
- (xvii) halogen,
- (xviii) C₁₋₄ alkoxy-carbonyl,
- (xix) C₆₋₁₀ aryloxy-carbonyl,
- (xx) C₃₋₆ cycloalkyloxy-carbonyl,
- (xxi) C₇₋₁₂ aralkyloxy-carbonyl,
- (xxii) C₁₋₅ alkanoyl,
- (xxiii) C₁₋₁₅ alkanoyloxy,
- (xxiv) carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, N-ethylcarbamoyl, N,N-diethylcarbamoyl, N-phenylcarbamoyl, pyrrolidinocarbamoyl, piperidinocarbamoyl, piperazinocarbamoyl, morpholinocarbamoyl or N-benzylcarbamoyl,

(xxv) N-methylcarbamoyloxy, N,N-dimethylcarbamoyloxy, N-ethylcarbamoyloxy, N-benzylcarbamoyloxy, N,N-dibenzylcarbamoyloxy or N-phenylcarbamoyloxy,

(xxvi) C₁₋₄ alkanoylamino,

(xxvii) C₆₋₁₀ arylcarbonylamino,

(xxviii) C₁₋₄ alkoxy-carbonylamino,

(xxix) C₇₋₁₂ aralkyloxycarbonyl,

(xxx) methanesulfonylamino, ethanesulfonylamino, butanesulfonylamino, benzenesulfonylamino, toluenesulfonylamino, naphthalenesulfonylamino, trifluoromethanesulfonylamino, 2-chloroethanesulfonylamino or 2,2,2-trifluoromethanesulfonylamino,

(xxxi) pyrrolidinyl, pyrrolyl, pyrazolyl, imidazolyl, furyl, thienyl, oxazolyl, isoxazolyl, isothiazolyl, thiazolyl, piperidinyl, pyridyl, piperazinyl, pyrimidinyl, pyranyl, tetrahydropyranyl, tetrahydrofuryl, indolyl, quinolyl, 1,3,4-oxadiazolyl, thieno[2,3-d]pyridyl, 1,2,3-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,3,4-triazolyl, tetrazolyl, 4,5-dihydro-1,3-dioxazolyl, tetrazolo[1,5-b]pyridazinyl, benzothiazolyl, benzoxazolyl, benzimidazolyl or benzothienyl,

(xxxii) heterocyclethio, heterocycleoxy, heterocycleamino or heterocyclecarbonylamino group which is derived by attachment of any of the heterocyclic groups (xxxi) defined above to the S, O, N atom or a carbonylamino group,

(xxxiii) di-C₁₋₄ alkylphosphinothioylamino,

(xxxiv) methoxyimino, ethoxyimino, 2-fluoroethoxyimino, carboxymethoxyimino, 1-carboxy-1-methylethoxyimino, 2,2,2-trichloroethoxycarbonylmethoxyimino, 1-(2,2,2-trichloroethoxycarbonyl)-1-methylethoxyimino, (2-aminothiazol-4-yl)methoxyimino or (1H-imidazol-4-yl)methoxyimino,

(xxxv) C₁₋₄ alkylsulfonyloxy,

(xxxvi) C₆₋₁₀ arylsulfonyloxy,

(xxxvii) di-C₆₋₁₀ arylphosphino-thioylamino,

(xxxviii) thiocarbamoylthio, N-methylthiocarbamoylthio, N,N-dimethylthiocarbamoylthio, N-ethylthiocarbamoylthio, N-benzylthiocarbamoylthio, N,N-dibenzylthiocarbamoylthio or N-phenylthiocarbamoylthio,

(xxxix) trimethylsilyloxy, t-butyldimethylsilyloxy, t-butyldiphenylsilyloxy or dimethylphenylsilyloxy,

(xL) trimethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl or dimethylphenylsilyl,

(xLi) C₁₋₄ alkylsulfinyl,

(xLii) C₆₋₁₀ arylsulfinyl,

(xLiii) C₁₋₄ alkylsulfonyl,

(xLiv) C₆₋₁₀ arylsulfonyl,

(xLv) C₁₋₄ alkoxy-carbonyloxy,

(xLvi) halo-C₁₋₄ alkyl,

(xLvii) halo-C₁₋₄ alkoxy, halo-C₁₋₄ alkylthio, halo-C₁₋₄ alkylsulfinyl or halo-C₁₋₄ alkylsulfonyl,


(xLviii) cyano, nitro, hydroxyl, carboxyl, sulfo, phosphono,

(xLix) C₁₋₄ alkylloxysulfonyl,

(L) C₆₋₁₀ aryloxysulfonyl,


(Li) C₇₋₁₂ aralkyloxysulfonyl, and

(Lii) di-C₁₋₄ alkyl oxyphosphoryl group, with the proviso that when R² is a hydrogen atom, R¹ is a group of the formula:




[wherein R^{3a} is hydrogen, C₁₋₄ alkyl, C₇₋₉ phenylalkyl or C₁₋₄ alkanoyl and R^{4a} is a hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, (di-C₁₋₄ alkylamino)-C₁₋₄ alkyl, tri-C₁₋₄ alkylsilyl-C₁₋₄ alkyl, C₂₋₄ alkenyl or pyridyl- or thiazolyl-C₁₋₂ alkyl wherein pyridyl or thiazolyl moiety may optionally be substituted with a halogen atom, or R^{3a} and R^{4a} taken together with the adjacent nitrogen atom constitute pyrrolidino) and A^o is pyridyl, pyrazinyl or thiazolyl, each of which may optionally be substituted with a halogen, C₁₋₄ alkyl, C₁₋₄ alkylthio or C₁₋₄ alkoxy),

and with the proviso that when



is $O_2N-CH=$;
 R^1 is



R^3 is hydrogen, C_1-5 alkyl or C_3-6 cycloalkyl;


R^4 is hydrogen, C_1-5 alkyl, C_3-6 cycloalkyl, benzyl or pyrimidinylmethyl; or

R^3 and R^4 together with the adjacent nitrogen atom constitute a cyclic amino group of pyrrolidinyl or piperazinyl; and

R^2 is hydrogen, C_1-5 alkyl or C_3-6 cycloalkyl,

A^0 is not a pyridyl substituted by C_1-4 haloalkyl, C_1-4 haloalkoxy, C_1-4 haloalkylthio, C_1-4 haloalkylsulfanyl, C_1-4 haloalkylsulfonyl, cyano, nitro or hydroxyl, or a salt thereof.

2. A compound as claimed in claim 1, wherein R^2 is hydrogen, R^1 is a group of the formula:



(wherein R^{3a} and R^{4a} are as defined in claim 1) and A^0 is heterocycle selected from the class consisting of pyridyl, pyrazinyl and thiazolyl, the said heterocycle mentioned just above for A^0 being optionally substituted with halogen, C_1-4 alkyl, C_1-4 alkylthio or C_1-4 alkoxy.

3. A compound as claimed in claim 1, wherein R^2 is other than hydrogen.

4. A compound as claimed in claim 1, wherein,

X^1 is nitro;

X^2 is hydrogen, C_1-2 alkoxy, carbonyl or C_1-2 alkylsulfonylthiocarbonyl;

R^1 is amino, mono- or di- C_1-4 alkylamino, halo- C_1-4 alkylamino, N- C_1-4 alkyl-N- C_1-2 alkanoylamino, N-halo- C_1-4 alkyl-N- C_1-2 alkanoylamino or C_1-2 alkanoylamino;

R^2 is hydrogen, C_1-2 alkoxy, di- C_1-2 alkylamino, C_1-4 alkyl, halo- C_1-4 alkyl or C_1-2 alkanoyl;

n is 0 or 1;

A^0 is 2- or 3-thienyl, 2- or 3-furyl, 2- or 3-pyrrolyl, 2-, 3- or 4-pyridyl, 2-, 4- or 5-oxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-imidazolyl, 3-, 4- or 5-isoxazolyl, 3-, 4- or 5-isothiazolyl, 3- or 5-(1,2,4-oxadiazolyl), 1,3,4-oxadiazolyl, 3- or 5-(1,2,4-thiadiazolyl), 1,3,4-thiadiazolyl, 4- or 5-(1,2,3-thiadiazolyl), 1,2,5-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1H- or 2H-tetrazolyl, N-oxido-2-, 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, N-oxido-2-, 4- or 5-pyrimidinyl, 3- or 4-pyridazinyl, pyrazinyl, N-oxido-3- or 4-pyridazinyl, benzofuryl, benzothiazolyl, benzoxazolyl, triazinyl, oxotriazinyl, tetrazolo[1,5-b]-pyridazinyl, triazolo[4,5-b]pyridazinyl, oxoimidazinyl, dioxotriazinyl, pyrrolidinyl, piperidinyl, pyranyl, thiopyranyl, 1,4-oxazinyl, morpholinyl, 1,4-thiazinyl, 1,3-thiazinyl, piperazinyl, benzimidazolyl, quinolyl, isoquinolyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, indoliziny, quinoliziny, 1,8-naphthyridinyl, purinyl, pteridinyl, dibenzofuranyl, carbazolyl, acridinyl, phenanthridinyl, phenazinyl, phenothiazinyl or phenoxyazinyl, each of which may optionally be substituted with halogen, C_1-4 alkyl, halo- C_1-4 alkyl, C_1-4 alkoxy, halo- C_1-4 alkoxy, C_1-4 alkylthio or halo- C_1-4 alkylthio or a salt thereof.

5. A compound as claimed in claim 1, wherein,

X^1 is nitro;

X^2 is hydrogen or C_{1-2} alkylsulfonylthiocarbamoyl;


R^1 is amino, mono- or di- C_{1-2} alkylamino, halo- C_{1-2} alkylamino, N- C_{1-2} alkyl-N- C_{1-2} alkanoylamino, N-halo- C_{1-2} alkyl-N- C_{1-2} alkanoylamino or C_{1-2} alkanoylamino;

R^2 is hydrogen, C_{1-2} alkoxy, di- C_{1-2} alkylamino, C_{1-4} alkyl, halo- C_{1-4} alkyl or C_{1-2} alkanoyl;

n is 1; and

A^0 is pyridyl, pyrazinyl or thiazolyl, each of which may optionally be substituted with halogen, C_{1-4} alkyl, halo- C_{1-4} alkyl, C_{1-4} alkoxy, halo- C_{1-4} alkoxy, C_{1-4} alkylthio or halo- C_{1-4} alkylthio or a salt thereof.

6. A compound as claimed in claim 1, of the formula



wherein:

X^{2a} is hydrogen, C_{1-4} alkoxycarbonyl or C_{1-4} alkylsulfonylthiocarbamoyl;


R^{2c} is hydrogen, C_{1-3} alkanoyl, C_{1-4} alkyl, mono- or di- C_{1-4} alkoxy- C_{1-4} alkyl, C_{7-9} aralkyl, mono- or di- C_{1-4} alkylamino or C_{1-4} alkoxy;

A^c is 3- or 4-pyridyl, pyrazinyl or 4- or 5-thiazolyl, each of which may optionally be substituted with halogen, C_{1-4} alkyl or C_{1-4} alkoxy;

n is 1; and

R^{3a} and R^{4a} are as defined in claim 1, or a salt thereof.

7. A compound as claimed in claim 1, which is a compound of the formula:



wherein:

X^{2a} is hydrogen, C_{1-4} alkoxycarbonyl or C_{1-4} alkylsulfonylthiocarbamoyl;


R^{1d} is amino, mono- or di- C_{1-4} alkylamino, N- C_{1-4} alkyl-N- C_{1-3} alkanoylamino, C_{7-9} aralkylamino, halogenothiazolyl- C_{1-2} alkylamino or C_{1-4} alkoxy- C_{1-2} alkylamino;

R^{2c} is hydrogen, C_{1-3} alkanoyl, C_{1-4} alkyl, mono- or di- C_{1-4} alkoxy- C_{1-4} alkyl, C_{7-9} aralkyl, mono- or di- C_{1-4} alkylamino or C_{1-4} alkoxy;

n is 0, 1 or 2; and

A^d is 3- or 4-pyridyl, pyrazinyl or 5-thiazolyl, each of which may optionally be substituted with halogen, C_{1-4} alkyl or C_{1-4} alkoxy, or a salt thereof.


8. A compound as claimed in claim 1, which is a compound of the formula:



wherein:

X^{2b} is hydrogen or C_{1-2} alkylsulfonylthiocarbamoyl;
 R^{1e} is amino, mono- or di- C_{1-2} alkylamino or N- C_{1-2} alkyl-N-formylamino;
 R^{2d} is hydrogen, C_{1-2} alkyl or C_{1-3} alkanoyl; and
 A^e is a group of the formula:


15



20

wherein Hal is a halogen atom, or a salt thereof.

25 9. A compound as claimed in claim 1, which is a compound of the formula:




30

wherein:

X^{2c} is hydrogen or methylsulfonylthiocarbamoyl;
 R^{1f} is amino, methylamino, dimethylamino or N-methyl-N-formylamino;
 R^{2d} is a hydrogen atom, formyl or C_{1-2} alkyl; and
 A^e is a group of the formula:

35




40

wherein Hal is a halogen atom, or a salt thereof.

10. A compound as claimed in claim 1, which is a compound of the formula:

50



55

wherein:

R^{1a} is amino, mono- or di-C₁₋₂ alkylamino or N-C₁₋₂ alkyl-N-formylamino;


R^{2a} is C₁₋₂ alkyl or formyl; and

Hal is a halogen atom, or a salt thereof.

5

11. A compound as claimed in claim 1, wherein the heterocycle is selected from the following group and being optionally substituted as defined in claim 1, the group consisting of 2- or 3-thienyl, 2- or 3-furyl, 2- or 3- pyrrolyl, 2-, 4- or 5-oxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-imidazolyl, 3-, 4- or 5-isoxazolyl, 3-, 4- or 5-isothiazolyl, 3- or 5-(1,2,4-oxadiazolyl), 1,3,4-oxadiazolyl, 3- or 5-(1,2,4-thiadiazolyl), 1,3,4-thiadiazolyl, 4- or 5-(1,2,3-thiadiazolyl), 1,2,5-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1H- or 2H-tetrazolyl, N-oxido-2-, 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, N-oxido-2-, 4- or 5-pyrimidinyl, 3- or 4-pyridazinyl, pyrazinyl, N-oxido-3- or 4-pyridazinyl, benzofuryl, benzothiazolyl, benzoxazolyl, triazinyl, oxotriazinyl, tetrazolo[1,5-b]pyridazinyl, trisolo[4,5-b]pyridazinyl, oxoimidazinyl, dioxotriazinyl, pyrrolidinyl, piperidinyl, pyranyl, thiopyranyl, 1,4-oxazinyl, morpholinyl, 1,4-thiazinyl, 1,3-thiazinyl, piperazinyl, benzimidazolyl, quinolyl, isoquinolyl, cinnolyl, phthalazinyl, quinazolinyl, quinoxalyl, indolizyl, quinolizyl, 1,8-naphthyridinyl, purinyl, pteridinyl, dibenzofuranyl, carbazolyl, acridinyl, phenanthridinyl, phenazinyl, phenothiazinyl and phenoxazinyl.
12. A compound as claimed in claim 1, selected from 1-[N-(6-chloro-3-pyridylmethyl)-N-methyl]amino-1-methylamino-2-nitroethylene, 1-(6-chloro-3-pyridylmethyl)amino-1-dimethylamino-2-nitroethylene, and 1-[N-(6-chloro-3-pyridylmethyl)-N-ethyl]amino-1-methylamino-2-nitroethylene.
13. An insecticidal/miticidal composition which comprises an insecticidal/miticidal effective amount of at least one of the α -unsaturated amines as claimed in any one of claims 1 to 12, or a salt thereof, together with a suitable carrier or carriers.
14. A process for preparing an α -unsaturated amine of the formula:


30



35

wherein the symbols are as defined in claim 1 or a salt thereof, which comprises
(1) reacting a compound of the formula:

40



45

or a salt thereof with a compound of the formula:

Y - W²

or a salt thereof, or

50

55

(2) reacting a compound of the formula:



or



or a salt thereof with a compound of the formula:



or a salt thereof, or
(3) reacting a compound of the formula:



or



(i) with a compound of the formula:



or a salt thereof, and then reacting the resulting product with a compound of the formula:



or a salt thereof, or (ii) with a compound of the formula:



or a salt thereof, and then reacting the resulting product with a compound of the formula:



or a salt thereof, or

10 (4) reacting a compound of the formula:



or a salt thereof with a compound of the formula:




or a salt thereof, or

25 (5) reacting a compound of the formula:




or a salt thereof with a compound of the formula:



or a salt thereof, or

(6) subjecting a compound of the formula:




45

or a salt thereof to hydrolysis reaction and then to decarboxylation reaction, or


50

55


(7) subjecting a compound of the formula:




or



or a salt thereof to alkylation, acylation, alkoxycarbonylation, sulfonylation or phosphorylation, in which formulas, R⁵ is a C₁₋₄ alkyl or aralkyl; when W¹ is



W² is R¹ and when W¹ is R¹, W² is



Y is a hydrogen atom or an alkali metal;

R³ is a hydrogen atom, alkyl, aryl, aralkyl, heterocyclic, acyl, alkoxycarbonyl, aryloxy, heterocycleoxy, arylsulfonyl, alkylsulfonyl, dialkoxylphosphoryl, alkoxy, hydroxyl, amino, dialkylamino, acylamino, alkoxycarbonylamino, alkylsulfonylamino, dialkoxylphosphorylamino, aralkyloxy or alkoxycarbonylalkyl; R⁴ is a hydrogen atom, or alkyl, cycloalkyl, alkenyl, cycloalkenyl or alkynyl which groups may optionally be substituted, or pyridyl- or thiazolyl-C₁₋₂ alkyl wherein pyridyl and thiazolyl moiety may optionally be substituted with a halogen atom; Hal is a halogen atom;

X³ is an electron-attracting group; R⁶ is a group attached through a nitrogen atom containing at least one hydrogen atom; and X¹, X², R¹, R², n and A⁰ are as defined in claim 1.


15. A method of combatting undesirable insects or mites, which comprises applying an insecticidal or miticidal effective amount of the compound of the formula [I⁰] defined in any one of claims 1 to 12 or a salt thereof to the said insects or mites or their habitat.

16. A method of claim 15, wherein the compound or salt is applied in a composition of the compound or salt with a suitable carrier or carriers.

17. A method of combatting undesirable insects or mites, which comprises applying an insecticidal or miticidal effective amount of the compound of the formula [I⁰] defined in claim 12.


Claims for the following Contracting State : ES

1. A process for preparing an α -unsaturated amine of the formula:




wherein:

one or X^1 and X^2 is an electron-attracting group and the other is a hydrogen atom or an electron-attracting group, wherein the said electron-attracting group is cyano, nitro, C_1 - 4 alkoxy carbonyl, carboxyl, C_6 - 10 aryloxy-carbonyl, heterocycleoxycarbonyl, C_1 - 4 alkylsulfonyl which may be substituted with halogen, aminosulfonyl, di- C_1 - 4 alkoxyphosphoryl, C_1 - 4 alkanoyl which may be substituted with halogen, C_1 - 4 alkylsulfonylthiocarbonyl, carbamoyl or halogen, or X^1 and X^2 together with the carbon atom to which they are attached form a ring of the formula:



R^1 is a group of the formula:




in which:

R^3 is hydrogen, C_1 - 20 alkyl, C_6 - 10 aryl, C_7 - 9 aralkyl, heterocycle, C_1 - 4 alkanoyl, C_6 - 10 aryl-carbonyl, C_1 - 4 alkoxy-carbonyl, C_6 - 10 aryloxy-carbonyl, heterocycleoxycarbonyl, C_6 - 10 arylsulfonyl, C_1 - 4 alkylsulfonyl, di- C_1 - 4 alkoxyphosphoryl, C_1 - 4 alkoxy, hydroxy, amino, di- C_1 - 4 alkylamino, C_1 - 4 alkanoylamino, C_1 - 4 alkoxy-carbonylamino, C_1 - 4 alkylsulfonylamino, di- C_1 - 4 alkoxyphosphorylamino, C_7 - 9 aralkyloxy or C_1 - 4 alkoxy-carbonyl- C_1 - 4 alkyl; and


R^4 is hydrogen, C_1 - 20 alkyl, C_3 - 6 cycloalkyl, C_2 - 6 alkenyl, C_3 - 6 cycloalkenyl or C_2 - 6 alkynyl, wherein each of the radicals defined for R^4 except for hydrogen may optionally be substituted by 1 to 3 substituents selected from the group consisting of hydroxy, C_1 - 4 alkoxy, halogen, di- C_1 - 4 alkylamino, C_1 - 4 alkylthio, C_1 - 3 alkanoylamino, C_1 - 4 alkylsulfonylamino, tri- C_1 - 4 alkylsilyl, pyridyl and thiazolyl, and each of the pyridyl and thiazolyl may further be substituted by halogen, or

R^3 and R^4 together with the adjacent nitrogen atom constitute a cyclic amino group of the formula:



R^2 is (1) hydrogen, (2) a group attached through a carbon atom selected from the class consisting of C_1 - 4 alkanoyl, C_1 - 20 alkyl, C_2 - 6 alkenyl, C_3 - 6 cycloalkyl, C_6 - 10 aryl, C_7 - 9 aralkyl and 3- or 4-pyridyl, the said group attached through a carbon atom being optionally substituted by 1 to 3 substituents selected from the class consisting of C_1 - 4 alkylthio, C_1 - 4 alkoxy, mono- or di- C_1 - 4 alkylamino, C_1 - 4 alkoxy-carbonyl, C_1 - 4 alkylsulfonyl, halogen and C_1 - 4 alkanoyl, (3) a group attached

through an oxygen atom selected from the class consisting of C₁₋₄ alkoxy, C₃₋₆ cycloalkoxy, C₂₋₄ alkenyloxy, C₃₋₆ cycloalkenyloxy, ethynyloxy, C₆₋₁₀ aryloxy, thienyloxy and hydroxy, the said group attached through an oxygen atom being optionally substituted by 1 to 3 substituents selected from the class consisting of halogen and phenyl, or (4) a group attached through a nitrogen atom of the formula:



wherein R³ and R⁴ have the meanings given above;

n is an integer of 0, 1 or 2;

A° is heterocycle,

wherein the heterocycle in the said heterocycle carbonyl for X¹ and X², the said heterocycle for R³, the heterocycle in the said heterocycleoxycarbonyl for R³,

and the said heterocycle for A° are a member selected from the class consisting of thienyl, furyl, pyrrolyl, pyridyl, oxazolyl, thiazolyl, pyrazolyl, imidazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, N-oxidopyridyl, pyrimidinyl, N-oxidopyrimidinyl, pyridazinyl, pyrazinyl, N-oxidopyridazinyl, benzofuryl, benzothiazolyl, benzoxazolyl, triazinyl, oxotriazinyl, tetrazo[1,5-b]-pyridazinyl, triazolo[4,5-b]pyridazinyl, oxoimidazinyl, dioxotriazinyl, pyrrolidinyl, piperidinyl, pyranyl, thiopyranyl, 1,4-oxazinyl, morpholinyl, 1,4-thiazinyl, 1,3-thiazinyl, piperazinyl, benzimidazolyl, quinolyl, isoquinolyl, indoliziny, quinoliziny, 1,8-naphthyridinyl, purinyl, pteridinyl, dibenzofuranyl, carbazolyl, acridinyl, phenanthridinyl, phenazinyl, phenothiazinyl and phenoxazinyl, the said heterocycle being optionally substituted by 1 to 5 substituents selected from the group consisting of,

(i) C₁₋₄ alkyl,

(ii) C₃₋₆ cycloalkyl,

(iii) C₆₋₁₀ aryl,

(iv) C₁₋₄ alkoxy,

(v) C₃₋₆ cycloalkyloxy,

(vi) C₆₋₁₀ aryloxy,

(vii) C₇₋₁₂ aralkyloxy

(viii) C₁₋₄ alkylthio,

(ix) C₃₋₆ cycloalkylthio,

(x) C₆₋₁₀ arylthio,

(xi) C₇₋₁₂ aralkylthio,

(xii) mono-C₁₋₄ alkylamino,

(xiii) di-C₁₋₄ alkylamino,

(xiv) C₃₋₆ cycloalkylamino,

(xv) C₆₋₁₀ arylamino,

(xvi) C₇₋₁₂ aralkylamino,

(xvii) halogen,

(xviii) C₁₋₄ alkoxy carbonyl,

(xix) C₆₋₁₀ aryloxy carbonyl,

(xx) C₃₋₆ cycloalkyloxy carbonyl,

(xxi) C₇₋₁₂ aralkyloxy carbonyl,

(xxii) C₁₋₅ alkanoyl,

(xxiii) C₁₋₁₅ alkanoyloxy,

(xxiv) carbamoyl, N-methylcarbamoyl, N,N-dimethylcarbamoyl, N-ethylcarbamoyl, N,N-diethylcarbamoyl, N-phenylcarbamoyl, pyrrolidinocarbamoyl, piperidinocarbamoyl, piperazinocarbamoyl, morpholinocarbamoyl or N-benzylcarbamoyl,

(xxv) N-methylcarbamoyloxy, N,N-dimethylcarbamoyloxy, N-ethylcarbamoyloxy, N-benzylcarbamoyloxy, N,N-dibenzylcarbamoyloxy or N-phenylcarbamoyloxy,

(xxvi) C₁₋₄ alkanoylamino,

(xxvii) C₆₋₁₀ arylcarbonylamino,

(xxviii) C₁₋₄ alkoxy carbonylamino,

(xxix) C₇₋₁₂ aralkyloxycarbonyl,

(xxx) methanesulfonylamino, ethanesulfonylamino, butanesulfonylamino, benzenesulfonylamino, toluenesulfonylamino, naphthalenesulfonylamino, trifluoromethanesulfonylamino, 2-chloroethanesulfonylamino or 2,2,2-trifluoromethanesulfonylamino,

(xxxi) pyrrolidinyl, pyrrolyl, pyrazolyl, imidazolyl, furyl, thienyl, oxazolyl, isoxazolyl, isothiazolyl, thiazolyl, piperidinyl, pyridyl, piperazinyl, pyrimidinyl, pyranyl, tetrahydropyranyl, tetrahydrofuryl, indolyl, quinolyl, 1,3,4-oxadiazolyl, thieno[2,3-d]pyridyl, 1,2,3-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,3,4-triazolyl, tetrazolyl, 4,5-dihydro-1,3-dioxazolyl, tetrazolo[1,5-b]-pyridazinyl, benzothiazolyl, benzoxazolyl, benzimidazolyl or benzothienyl,

(xxxii) heterocyclethio, heterocycleoxy, heterocycleamino or heterocyclecarbonylamino group which is derived by attachment of any of the heterocyclic groups (xxxi) defined above to the S, O, N atom or a carbonylamino group,

(xxxiii) di-C₁₋₄ alkylphosphinothioylamino,

(xxxiv) methoxyimino, ethoxyimino, 2-fluoroethoxyimino, carboxymethoxyimino, 1-carboxy-1-methylethoxyimino, 2,2,2-trichloroethoxycarbonylmethoxyimino, 1-(2,2,2-trichloroethoxycarbonyl)-1-methylethoxyimino, (2-aminothiazol-4-yl)methoxylmino or (1H-imidazol-4-yl)methoxyimino,

(xxxv) C₁₋₄ alkylsulfonyloxy,

(xxxvi) C₆₋₁₀ arylsulfonyloxy,

(xxxvii) di-C₆₋₁₀ arylphosphino-thioylamino,

(xxxviii) thiocarbamoylthio, N-methylthiocarbamoylthio, N,N-dimethylthiocarbamoylthio, N-ethylthiocarbamoylthio, N-benzylthiocarbamoylthio, N,N-dibenzylthiocarbamoylthio or N-phenylthiocarbamoylthio,

(xxxix) trimethylsilyloxy, t-butyldimethylsilyloxy, t-butyldiphenylsilyloxy or dimethylphenylsilyloxy,

(xL) trimethylsilyl, t-butyldimethylsilyl, t-butyldiphenylsilyl or dimethylphenylsilyl,

(xLi) C₁₋₄ alkylsulfinyl,

(xLii) C₆₋₁₀ arylsulfinyl,

(xLiii) C₁₋₄ alkylsulfonyl,

(xLiv) C₆₋₁₀ arylsulfonyl,

(xLv) C₁₋₄ alkoxy-carbonyloxy,

(xLvi) halo-C₁₋₄ alkyl,

(xLvii) halo-C₁₋₄ alkoxy, halo-C₁₋₄ alkylthio, halo-C₁₋₄ alkylsulfinyl or halo-C₁₋₄ alkylsulfonyl,


(xLviii) cyano, nitro, hydroxyl, carboxyl, sulfo, phosphono,

(xLix) C₁₋₄ alkyloxysulfonyl,


(L) C₆₋₁₀ aryloxysulfonyl,

(Li) C₇₋₁₂ aralkyloxysulfonyl, and

(Lii) di-C₁₋₄ alkyloxyphosphoryl group, with the proviso that when R² is a hydrogen atom, R¹ is a group of the formula,




[wherein R^{3a} is hydrogen, C₁₋₄ alkyl, C₇₋₉ phenylalkyl or C₁₋₄ alkanoyl and R^{4a} is a hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy-C₁₋₄ alkyl, (di-C₁₋₄ alkylamino)-C₁₋₄ alkyl, tri-C₁₋₄ alkylsilyl-C₁₋₄ alkyl, C₂₋₄ alkenyl or pyridyl- or thiazolyl-C₁₋₂ alkyl wherein pyridyl or thiazolyl moiety may optionally be substituted with a halogen atom, or R^{3a} and R^{4a} taken together with the adjacent nitrogen atom constitute pyrrolidino] and A^o is pyridyl, pyrazinyl or thiazolyl, each of which may optionally be substituted with a halogen, C₁₋₄ alkyl, C₁₋₄ alkylthio or C₁₋₄ alkoxy), and with the proviso that when



is $O_2N-CH=$;
 R^1 is


5



R^3 is hydrogen, C_{1-5} alkyl or C_{3-6} cycloalkyl;
 R^4 is hydrogen, C_{1-5} alkyl, C_{3-6} cycloalkyl, benzyl or pyrimidinylmethyl; or
 R^3 and R^4 together with the adjacent nitrogen atom constitute a cyclic amino group of pyrrolidinyl or piperazinyl; and
 R^2 is hydrogen, C_{1-5} alkyl or C_{3-6} cycloalkyl,
 A^0 is not a pyridyl substituted by C_{1-4} haloalkyl, C_{1-4} haloalkoxy,
 C_{1-4} haloalkylthio, C_{1-4} haloalkylsulfinyl, C_{1-4} haloalkylsulfonyl, cyano, nitro or hydroxyl,
 or a salt thereof,
 which comprises

(1) reacting a compound of the formula:

20



25

or a salt thereof with a compound of the formula:


$Y - W^2$

30


or a salt thereof, or

(2) reacting a compound of the formula:

35




40



45


or

50



or a salt thereof with a compound of the formula:


55



5

or a salt thereof, or
(3) reacting a compound of the formula:

10




15

or



(i) with a compound of the formula:

20



25

or a salt thereof, and then reacting the resulting product with a compound of the formula:



30


or a salt thereof, or (ii) with a compound of the formula:



35


or a salt thereof, and then reacting the resulting product with a compound of the formula:

40



or a salt thereof, or
(4) reacting a compound of the formula:

45



50


or a salt thereof with a compound of the formula:




55

or a salt thereof, or

(5) reacting a compound of the formula:




or a salt thereof with a compound of the formula:




or a salt thereof, or

(6) subjecting a compound of the formula:




or a salt thereof to hydrolysis reaction and then to decarboxylation reaction, or


(7) subjecting a compound of the formula:




or



or a salt thereof to alkylation, acylation, alkoxycarbonylation, sulfonylation or phosphorylation, in which formulas, R^5 is a C_1-4 alkyl or aralkyl; when W^1 is



W^2 is R^1 and when W^1 is R^1 , W^2 is




Y is a hydrogen atom or an alkali metal;

R^3 is a hydrogen atom, alkyl, aryl, aralkyl, heterocyclic, acyl, alkoxycarbonyl, aryloxy carbonyl,

heterocycleoxycarbonyl, arylsulfonyl, alkylsulfonyl, dialkoxyphosphoryl, alkoxy, hydroxyl, amino, dialkylamino, acylamino, alkoxy-carbonylamino, alkylsulfonylamino, dialkoxyphosphorylamino, aralkyloxy or alkoxy-carbonylalkyl; R⁴ is a hydrogen atom, or alkyl, cycloalkyl, alkenyl, cycloalkenyl or alkynyl which groups may optionally be substituted, or pyridyl- or thiazolyl-C₁₋₂ alkyl wherein pyridyl and thiazolyl moiety may optionally be substituted with a halogen atom; Hal is a halogen atom; X³ is an electron-attracting group; R⁶ is a group attached through a nitrogen atom containing at least one hydrogen atom; and X¹, X², R¹, R², n and A⁰ are defined as above.

2. A process as claimed in claim 1, wherein R² is hydrogen, R¹ is a group of the formula:



(wherein R^{3a} and R^{4a} are as defined in claim 1) and A⁰ is heterocycle selected from the class consisting of pyridyl, pyrazinyl and thiazolyl, the said heterocycle mentioned just above for A⁰ being optionally substituted with halogen, C₁₋₄ alkyl, C₁₋₄ alkylthio or C₁₋₄ alkoxy.

3. A process as claimed in claim 1, wherein R² is other than hydrogen.

4. A process as claimed in claim 1, wherein:

X¹ is nitro;

X² is hydrogen, C₁₋₂ alkoxy-carbonyl or C₁₋₂ alkylsulfonylthiocarbonyl;

R¹ is amino, mono- or di-C₁₋₄ alkylamino, halo-C₁₋₄ alkylamino, N-C₁₋₄ alkyl-N-C₁₋₂ alkanoylamino, N-halo-C₁₋₄ alkyl-N-C₁₋₂ alkanoylamino or C₁₋₂ alkanoylamino;

R² is hydrogen, C₁₋₂ alkoxy, di-C₁₋₂ alkylamino, C₁₋₄ alkyl, halo-C₁₋₄ alkyl or C₁₋₂ alkanoyl;

n is 0 or 1;

A⁰ is 2- or 3-thienyl, 2- or 3-furyl, 2- or 3-pyrrolyl, 2-, 3- or 4-pyridyl, 2-, 4- or 5-oxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-imidazolyl, 3-, 4- or 5-isoxazolyl, 3-, 4- or 5-isothiazolyl, 3- or 5-(1,2,4-oxadiazolyl), 1,3,4-oxadiazolyl, 3- or 5-(1,2,4-thiadiazolyl), 1,3,4-thiadiazolyl, 4- or 5-(1,2,3-thiadiazolyl), 1,2,5-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1H- or 2H-tetrazolyl, N-oxido-2-, 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, N-oxido-2-, 4- or 5-pyrimidinyl, 3- or 4-pyridazinyl, pyrazinyl, N-oxido-3- or 4-pyridazinyl, benzofuryl, benzothiazolyl, benzoxazolyl, triazinyl, oxotriazinyl, tetrazolo[1,5-b]-pyridazinyl, triazolo[4,5-b]pyridazinyl, oxoimidazinyl, dioxotriazinyl, pyrrolidinyl, piperidinyl, pyranyl, thiopyranyl, 1,4-oxazinyl, morpholinyl, 1,4-thiazinyl, 1,3-thiazinyl, piperazinyl, benzimidazolyl, quinolyl, isoquinolyl, cinnolyl, phthalazinyl, quinazolinyl, quinoxalinyl, indolizynyl, quinolizynyl, 1,8-naphthyridinyl, purinyl, pteridinyl, dibenzofuran, carbazolyl, acridinyl, phenanthridinyl, phenazinyl, phenothiazinyl or phenoxyazinyl, each of which may optionally be substituted with halogen, C₁₋₄ alkyl, halo-C₁₋₄ alkyl, C₁₋₄ alkoxy, halo-C₁₋₄ alkoxy, C₁₋₄ alkylthio or halo-C₁₋₄ alkylthio or a salt thereof.

5. A process as claimed in claim 1, wherein:

X¹ is nitro;

X² is hydrogen or C₁₋₂ alkylsulfonylthiocarbonyl;


R¹ is amino, mono- or di-C₁₋₂ alkylamino, halo-C₁₋₂ alkylamino, N-C₁₋₂ alkyl-N-C₁₋₂ alkanoylamino, N-halo-C₁₋₂ alkyl-N-C₁₋₂ alkanoylamino or C₁₋₂ alkanoylamino;

R² is hydrogen, C₁₋₂ alkoxy, di-C₁₋₂ alkylamino, C₁₋₄ alkyl, halo-C₁₋₄ alkyl or C₁₋₂ alkanoyl;

n is 1; and

A⁰ is pyridyl, pyrazinyl or thiazolyl, each of which may optionally be substituted with halogen, C₁₋₄ alkyl, halo-C₁₋₄ alkyl, C₁₋₄ alkoxy, halo-C₁₋₄ alkoxy, C₁₋₄ alkylthio or halo-C₁₋₄ alkylthio or a salt thereof.


6. A process as claimed in claim 1 for preparing a compound of the formula



wherein:

- X^{2a} is hydrogen, C_1 - 4 alkoxy carbonyl or C_1 - 4 alkylsulfonylthiocarbamoyl;
 R^{2c} is hydrogen, C_1 - 3 alkanoyl, C_1 - 4 alkyl, mono- or di- C_1 - 4 alkoxy- C_1 - 4 alkyl, C_7 - 9 aralkyl, mono- or di- C_1 - 4 alkylamino or C_1 - 4 alkoxy;
 A^c is 3- or 4-pyridyl, pyrazinyl or 4- or 5-thiazolyl, each of which may optionally be substituted with halogen, C_1 - 4 alkyl or C_1 - 4 alkoxy;
 n is 1; and
 R^{3a} and R^{4a} are as defined in claim 1, or a salt thereof.


7. A process as claimed in claim 1 for preparing a compound of the formula:



wherein:

- X^{2a} is hydrogen, C_1 - 4 alkoxy carbonyl or C_1 - 4 alkylsulfonylthiocarbamoyl;
 R^{1d} is amino, mono- or di- C_1 - 4 alkylamino, N - C_1 - 4 alkyl- N - C_1 - 3 alkanoylamino C_7 - 9 aralkylamino, halogenothiazolyl- C_1 - 2 alkylamino or C_1 - 4 alkoxy- C_1 - 2 alkylamino;
 R^{2c} is hydrogen, C_1 - 3 alkanoyl, C_1 - 4 alkyl, mono- or di- C_1 - 4 alkoxy- C_1 - 4 alkyl, C_7 - 9 aralkyl, mono- or di- C_1 - 4 alkylamino or C_1 - 4 alkoxy;
 n is 0, 1 or 2; and
 A^d is 3- or 4-pyridyl, pyrazinyl or 5-thiazolyl, each of which may optionally be substituted with halogen, C_1 - 4 alkyl or C_1 - 4 alkoxy, or a salt thereof.

8. A process as claimed in claim 1 for preparing a compound of the formula:




wherein:

- X^{2b} is hydrogen or C_1 - 2 alkylsulfonylthiocarbamoyl;
 R^{1e} is amino, mono- or di- C_1 - 2 alkylamino or N - C_1 - 2 alkyl- N -formylamino;
 R^{2d} is hydrogen, C_1 - 2 alkyl or C_1 - 3 alkanoyl; and
 A^e is a group of the formula:



wherein Hal is a halogen atom, or a salt thereof.

- 10 9. A process as claimed in claim 1 for preparing a compound of the formula:




wherein:

- 20 X^{2c} is hydrogen or methylsulfonylthiocarbamoyl;
 R^{1f} is amino, methylamino, dimethylamino or N-methyl-N-formylamino;
 R^{2d} is a hydrogen atom, formyl or C₁₋₂ alkyl; and
 A^e is a group of the formula:



wherein Hal is a halogen atom, or a salt thereof.

- 35 10. A process as claimed in claim 1 for preparing a compound of the formula:



wherein:

- 45 R^{1e} is amino, mono- or di-C₁₋₂ alkylamino or N-C₁₋₂ alkyl-N-formylamino;
 R^{2e} is C₁₋₂ alkyl or formyl; and
 Hal is a halogen atom, or a salt thereof.

- 50 11. A process as claimed in claim 1, wherein the heterocycle is selected from the following group and being optionally substituted as defined in claim 1, the group consisting of 2- or 3-thienyl, 2- or 3-furyl, 2- or 3-pyrrolyl, 2-, 4- or 5-oxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-imidazolyl, 3-, 4- or 5-isoxazolyl, 3-, 4- or 5-isothiazolyl, 3- or 5-(1,2,4-oxadiazolyl), 1,3,4-oxadiazolyl, 3- or 5-(1,2,4-thiadiazolyl), 1,3,4-thiadiazolyl, 4- or 5-(1,2,3-thiadiazolyl), 1,2,5-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1H- or 2H-tetrazolyl, N-oxido-2-, 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, N-oxido-2-, 4- or 5-pyrimidinyl, 3- or 4-pyridazinyl, pyrazinyl, N-oxido-3- or 4-pyridazinyl, benzofuryl, benzothiazolyl, benzoxazolyl, triazinyl, oxotriazinyl, tetrazolo[1,5-b]pyridazinyl, triazolo[4,5-b]pyridazinyl, oxoimidazinyl, dioxotriazinyl, pyrrolidinyl, pyranyl, thiopyranyl, 1,4-oxazinyl, morpholinyl, 1,4-thiazinyl, 1,3-thiazinyl, piperazinyl, benzimidazolyl, quinolyl, isoquinolyl, cinnolyl, phthalazinyl, quinazolinyl, quinox-
- 55


alanyl, indoliziny, quinoliziny, 1,8-naphthyridiny, puriny, pteridiny, dibenzofuranyl, carbazolyl, acridiny, phenanthridiny, phenaziny, phenothiaziny and phenoxaziny.

12. A process as claimed in claim 1 for the preparation of a compound selected from 1-[N-(6-chloro-3-pyridylmethyl)-N-methyl]amino-1-methylamino-2-nitroethylene, 1-(6-chloro-3-pyridylmethyl)amino-1-dimethylamino-2-nitroethylene, and 1-[N-(6-chloro-3-pyridylmethyl)-N-ethyl]amino-1-methylamino-2-nitroethylene.
13. A process for preparing an insecticidal/miticidal composition which comprises mixing an insecticidal/miticidal effective amount of at least one of the α -unsaturated amines as prepared according to any one of claims 1 to 12, or a salt thereof, together with a suitable carrier or carriers.
14. A method of combatting undesirable insects or mites, which comprises applying an insecticidal or miticidal effective amount of the compound of the formula [I°] prepared according to any one of claims 1 to 12 or a salt thereof to the said insects or mites or their habitat.
15. A method of claim 14, wherein the compound or salt is applied in a composition of the compound or salt with a suitable carrier or carriers.

Patentansprüche


Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. α -Ungesättigtes Amin der Formel



worin

eines von X^1 und X^2 eine elektronenanziehende Gruppe ist und das andere ein Wasserstoffatom oder eine elektronenanziehende Gruppe ist, in welcher die elektronenanziehende Gruppe Cyano, Nitro, C_1 -4-Alkoxycarbonyl, Carboxy, C_6 -10-Aryloxycarbonyl, Heterocyclyloxycarbonyl, C_1 -4-Alkylsulfonyl, welches mit Halogen substituiert sein kann, Aminosulfonyl, Di- C_1 -4-alkoxyphosphoryl, C_1 -4-Alkanoyl, welches mit Halogen substituiert sein kann, C_1 -4-Alkylsulfonylthiocarbamoyl, Carbamoyl oder Halogen ist, oder X^1 und X^2 zusammen mit dem Kohlenstoffatom, an welches sie gebunden sind, einen Ring der Formel



bilden;

R^1 eine Gruppe der Formel

